



# WeedMaster®

## Depend on Weedmaster®.

For years, growers and ranchers have counted on Weedmaster® for consistent weed control in sugarcane, pastures, small grains and fallowland. Weedmaster is a selective, post-emerge herbicide that provides tough, long-lasting control of a broad spectrum of annual, biennial and perennial broadleaf weeds. The growth regulating effects of 2,4-D and dicamba interfere with the normal development of weeds, ultimately causing plant death.

### CONSISTENT CONTROL IN A CONCENTRATED FORMULA

- Achieve outstanding control of brush and broadleaf weeds in a concentrated formula
- Controls tough weeds such as horsenettle, thistles and blackberry
- Safe for desirable grasses
- Pre-emergence control with residual activity

<b>EPA REG. NO.</b>	71368-34
<b>ACTIVE INGR.</b>	dicamba (12.4%), 2,4-D (35.7%)
<b>FORMULATION</b>	liquid
<b>CHEM. FAMILY</b>	benzoic acid, phenoxy-carboxylic acid
<b>HRAC NO.</b>	4
<b>MODE OF ACT.</b>	auxin growth regulator
<b>REI</b>	48 hours
<b>SIGNAL WORD</b>	danger
<b>RESTRICTED</b>	no
<b>PACKAGE SIZE</b>	5 x 1 gal, 2 x 2.5 gal, 30 gal, 265 gal tote, bulk



**Nufarm**

Grow a better tomorrow.



### KEY USES

Conservation Reserve Program (CRP) land  
 Fallowland  
 General farmstead  
 Sorghum  
 Grass (*hay or silage*)  
 Pasture  
 Rangeland  
 Sugarcane  
 Wheat

### KEY WEEDS CONTROLLED

(See product label for complete list)

#### ANNUALS

Buckwheat, wild  
 Burdock  
 Buttercup  
 Cockle, cow  
 Cocklebur, common  
 Dogfennel  
 Evening primrose  
 Fleabane, annual  
 Flixweed  
 Knotweed  
 Kochia  
 Lambsquarters, common  
 Maretail (*horseweed*)  
 Morningglory; ivyleaf, tall  
 Mustards; annual, tansy  
 Pigweed; prostrate, redroot, smooth, tumble  
 Purslane, common  
 Ragweed; common, lanceleaf, western  
 Shepherd's-purse  
 Smartweed, Pennsylvania  
 Sneezeweed, bitter sowthistle  
 Thistle, Russian  
 Velvetleaf

#### BIENNIALS & PERENNIALS

Bindweed, field  
 Bittercress  
 Chicory  
 Dandelion, common  
 Dock, curly  
 Goldenrod, Missouri  
 Groundsel, Texas  
 Honeysuckle, hairy  
 Horsenettle, Carolina  
 Ivy, poison  
 Marshelder  
 Mesquite  
 Persimmon, eastern  
 Prickly lettuce  
 Smartweed  
 Sorrel, red  
 Thistle, bull  
 Vetch, hairy

### USE SUMMARY

#### RATES

0.5 pints per acre to 4 to 5 pints per acre for control of annual, biennial and perennial weeds. See label for per acre rate according to weed growth stage.

Apply to actively growing weeds as aerial, broadcast, band, or spot spray applications.

Apply with water or sprayable fluid fertilizer as a carrier. Sprayable fluid fertilizer may be used as the carrier in preplant or pre-emergence uses. Post-emergence uses with sprayable fluid fertilizer may be made only on pasture, hayland, or wheat crops.

Surfactants and/or low use rate liquid fertilizers (28-0-0, 32-0-0) or crop oil concentrate may be used with this product to improve burndown of emerged weeds. (Do not use crop oil concentrate for feed crops – refer to crop oil concentrate label for complete directions.)

**Aerial Application:** Use 3 to 10 gallons of water per acre.

**Ground Application (Broadcast):** Use 5 to 40 gallons per broadcast acre.

*See label for complete directions for use.*

