

**RESTRICTED-USE PESTICIDE  
DUE TO HAZARD TO NONTARGET ORGANISMS**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

**PELLETED  
BAIT**



Photo is not representative of a label placement

**P.C.Q.<sup>®</sup> PRO**

**KILLS CALIFORNIA GROUND SQUIRRELS,  
MEADOW & PINE VOLES, NORWAY RATS,  
ROOF RATS & HOUSE MICE**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panels for First Aid and additional precautionary statements.

ACTIVE INGREDIENT:

Diphacinone (CAS #82-66-6): . . . . . 0.005%

OTHER INGREDIENTS: . . . . . 99.995%  
100.000%

**NET WT:**

**12 lbs (5.4 kg)**

#### PRECAUTIONARY STATEMENTS

##### Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed. Keep away from children, domestic animals and pets.

All handlers (including applicators) must wear shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

##### User Safety Requirements

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

##### FIRST AID

HAVE THE PRODUCT CONTAINER OR LABEL WITH YOU WHEN CALLING A POISON CONTROL CENTER OR DOCTOR OR GOING FOR TREATMENT

If Swallowed:

- Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.
- Have a person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

##### NOTE TO PHYSICIAN

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

##### TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or 1-877-854-2494 at once.

##### NOTE TO VETERINARIAN

**Anticoagulant Diphacinone:** For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K<sub>1</sub>.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for sites, pests, and application methods described on this label.

Do not sell this product in individual containers holding less than 4 pounds of bait.

##### USE RESTRICTIONS – For All Uses

Use only in rodent infested areas where exposure to agricultural crops will not occur. Do not contaminate food or food stuffs. Do not apply this bait at sites or to control pests not indicated on this label. Do not apply this product by application methods not specified in this label. Do not pile bait. Do not apply bait in or near actively growing food or feed crops or in areas where contaminating food or surfaces that may contact food is likely. Do not apply bait by aircraft at sites where bait may contaminate bodies of water. When applying bait along canals, and other waterways, use equipment that directs bait away from the water.

##### NORWAY RATS ROOF RATS AND HOUSE MICE

**IMPORTANT:** Do not expose children, pets, or nontarget animals to rodenticides. To prevent accidents:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

**USE RESTRICTIONS:** This product can only be used to control Norway rats, roof rats and house mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

Tamper-resistant bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

In active rodent burrows, apply this product to control Norway rats, and roof rats within or beyond 100 feet of buildings and man-made structures, provided that infestations of these rodents have been confirmed. Bait must be placed no less than 6 inches into active Norway/roof rat burrows. Do not broadcast bait. Because Norway/roof rat infestations may occur in areas farther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove food trash, garbage, clutter and debris.

##### APPLICATION DIRECTIONS:

**Norway and Roof Rats** - Apply 4 to 16 oz. of bait (usually at intervals of 15 to 30 feet) per placement in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease. Apply 1 to 2 oz. of bait deep into active rat burrows.

**House Mice:** Apply ¼ to ½ oz. (1 to 2 level tablespoons) of bait at 8 to 12 foot intervals in infested areas. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Using waterproof gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

## RESTRICTED-USE PESTICIDE DUE TO HAZARD TO NONTARGET ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

# P.C.Q.® PRO

##### ACTIVE INGREDIENT:

Diphacinone (CAS #62-66-6) ..... 0.005%

OTHER INGREDIENTS: ..... 99.995%

TOTAL 100.000%

## KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for First Aid and additional precautionary statements.

##### ENVIRONMENTAL HAZARDS

This product is extremely toxic to birds, fish and other wildlife. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

##### ENDANGERED SPECIES CONSIDERATIONS

**NOTICE:** It is a Federal offense to use any pesticide product in a manner that results in the death of a member of an endangered species. The use of this product may pose a hazard to Federally designated endangered/threatened species. Suggested measures to protect endangered species in your area may exist in an EPA Interim Measures pamphlet. You may call EPA's Endangered Species Hotline, 1-800-477-3813, to find out if an Interim Measures pamphlet exists in your area and have one sent to you. You can also consult with your local area extension office or pesticide lead agency to determine whether they have imposed any requirements in your area to protect endangered and threatened species.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. **[Plastic:]** Triple rinse (or equivalent), then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

## NET WEIGHT: 12 lbs (5.45 kg)

EPA REG. NO.: 12455-136

EPA EST. NO.: 12455-WI-1

Manufactured by:



3699 Kinsman Blvd.  
Madison, WI 53704 U.S.A.  
www.belllabs.com

MADE IN USA

#### CALIFORNIA GROUND SQUIRRELS

**USE RESTRICTIONS:** For control of California Ground Squirrels (*Spermophilus beecheyi*) in industrial, commercial, and residential areas. Product must be used in a secured, tamper-resistant bait station, or as a spot baiting application at the following sites: during the dormant season only for vineyards, fruit tree orchards (stone and pome), nut tree orchards, avocados, and citrus groves; rangeland, and pastures; noncrop borders of the listed use sites as follows: outside of fence rows, rights-of-way, canal banks, ditch banks, highways, levees, railroad lines, utilities. It may also be used in and around man-made structures such as agricultural buildings (but not inside of pens), homes, other permanent or temporary structures; campgrounds, golf courses, parks, athletic fields, recreational areas; ornamental plants; lawns; commercial horticultural nurseries, and plantations of forest trees.

Do not graze livestock or plant food or feed crops in spot-treated areas while bait is present. Applications in orchards, groves, and vineyards may only be made after harvest and during the dormant period and may not be made after tree and vine growth resumes in the spring.

Do not apply by aerial application. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers can be in the area during application.

**BAIT STATIONS:** Place secured tamper-resistant bait stations where California Ground Squirrel activity is present, usually near active burrows. Place bait stations up to 75 feet apart or near active burrows. Apply 2–4 lb. of bait per bait station. Bait stations should be checked every 3–5 days and filled as necessary until signs of bait consumption has ceased.

**SPOT BAITING:** Evenly scatter 1/4 cup in a measuring cup (0.1 lb.) of bait no larger than 40–50 square feet (i.e. a circular area no more than 8 feet in diameter), around the active California Ground Squirrel burrows and runways. Do not over-bait. Do not place bait in piles. Using the same procedure, make a second application 4 days after the first. Do not apply more than 10 pounds per acre per year.

**Surveillance and Follow-Up:** Dead ground squirrels should begin to appear in or near treated areas within 4 to 5 days after the first bait application. Visit the application site to monitor the effects of treatment 4 days after the first application and at subsequent intervals of 1 to 2 days. Wearing gloves, collect and properly dispose of visible carcasses of ground squirrels and other species found in or near treated areas. Dead animals buried on site must be buried a minimum of 18 inches below the ground surface, preferably deeper. Continue to collect and dispose of dead ground squirrels and search for non-target animals for at least 2 weeks after the last bait application, or longer if carcasses are still being found at that time. Carcass collection and burial should be performed in the late afternoon, near sundown, to reduce the potential for nocturnal scavengers or predators to find carcasses on the ground surface. Wear gloves and use leakproof plastic bags or other suitable containers for transporting carcasses not buried on site. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

#### MEADOW AND PINE VOLES

##### INDUSTRIAL, COMMERCIAL, AND RESIDENTIAL AREAS

**USE RESTRICTIONS:** For control of meadow voles (*Microtus pennsylvanicus*) and pine voles (*Microtus pinetorum*) in industrial, commercial, and residential areas. Areas include: in and around man-made structures such as agricultural buildings (do not apply in animal pens), homes & lawns (hand baiting into the burrow or tamper resistant bait station must be used for all residential uses and within 100 feet of a home), other permanent or temporary structures; campgrounds; golf courses; ornamental plants; commercial nurseries. If placed into a tamper-resistant bait station, may also be used in parks, athletic fields, and recreational areas.

Do not apply by aerial application. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers can be in the area during application.

**BURROW BAITING:** Identify surface trails and the mouths of holes leading to underground burrow systems. Place 1½ oz. (2 tablespoons) into each active burrow. Do not allow bait to be exposed on the surface. Do not disturb the runway system.

**BAIT STATIONS:** Place 2 lb. of bait in a tamper-resistant bait station in areas frequented by voles (near surface trails and burrows).

**GROUND BROADCAST BAITING:** Broadcast by hand or commercial spreader only within 100 feet of a man-made structure and only in commercial nurseries. Wear rubber gloves during hand-broadcast operations. Apply 10 lb. per acre. For high population areas, make a second application (10 lb. per acre) one to two months later. For pine vole application in states east of the Mississippi River, a rate of 20 lb. per acre may be required. Do not exceed 20 lb. per acre per calendar year.

#### FRUIT TREE ORCHARDS (DORMANT)

**USE RESTRICTIONS:** For control of meadow voles (*Microtus pennsylvanicus*) and pine voles (*M. pinetorum*) in pome fruit and stone fruit tree orchards in the states listed below\*. Aerial application is only permitted in those following states where aerial is listed next to the state name. Apply only after fall harvest (including drops), before new spring growth and when no rain or snow is expected for three consecutive days. Do not graze animals in treated areas. **California**, **Connecticut**, **Georgia**, **Idaho (aerial)**, **Massachusetts**, **Michigan (aerial)**, **Missouri**, **Montana (aerial)**, **North Carolina**, **New Hampshire**, **Ohio (aerial)**, **Oregon (aerial)**, **Pennsylvania (aerial)**, **South Carolina**, **Utah (aerial)**, **Virginia (aerial)**, **Vermont (aerial)**, **Washington (aerial)**, and **West Virginia (aerial)**.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers can be in the area during application.

**BURROW BAITING:** Identify surface trails and the mouths of holes leading to underground burrow systems. Place 1½ oz. (2 tablespoons) into each active burrow. Do not allow bait to be exposed on the surface. Do not disturb the runway system.

**BAIT STATIONS:** Place 2 lb. of bait in a tamper-resistant bait station in areas frequented by voles (near surface trails and burrows).

**BROADCAST BAITING:** Broadcast by hand, commercial spreader, or aerial (see use restrictions). Wear rubber gloves during hand-broadcast operations. Apply 10 lb. per acre. For high population areas, make a second application (10 lb. per acre) one to two months later.



Product Code: PQ1012

062916/05-16