

TriStar[®] 8.5 SL



Fast Knockdown in a Lasting Foliar Spray.

TriStar[®] 8.5 SL provides excellent, proven control of a wide range of outdoor ornamental insect pests

QUICK KNOCKDOWN AND LASTING RESIDUAL CONTROL

- Easy to mix and measure small volume applications
- Compatible with many commonly used surfactants, miticides and insecticides
- Contact, systemic and translaminar activity
- Fast-acting, broad-spectrum, IPM compatible
- Complete life cycle control from egg to adult on select insects
- Stable in sunlight
- Long-lasting residual control

EPA REG. NO.	8033-106-1001
ACTIVE INGR.	acetamiprid (8.5% by wt.)
FORMULATION	soluble liquid
CHEM. FAMILY	neonicotinoid
IRAC NO.	4A
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	8 x 4 oz tip-n-pour, 4 x 1 qt, 4 x 1 gal

 RESIDENTIAL

 COMMERCIAL

 LANDSCAPE ORNAMENTAL



Grow a better tomorrow.

TriStar[®] 8.5 SL

KEY USES

Annual and perennial flowering plants
Ornamental plants
Shrubs
Vines

KEY PESTS CONTROLLED

(See product label for complete list)

Adelgids
Aphids
Citrus thrips
Crane fly larvae
European pine sawflies
Hard and soft scales
Leaf eating beetles
Leafhoppers
Leafminers
Mealybugs
Plant bugs
Psyllids
Strawberry weevils
Thrips
Whiteflies

USE SUMMARY

For control of insect pests, apply either as foliar broadcast spray to obtain thorough and uniform spray coverage of the plants or via a basal bark or injection treatment. For foliar broadcast sprays, choose a finished spray volume appropriate for the size of the plants and amount of foliage which will provide thorough coverage throughout the canopy. For optimum control, allow at least 6 hours before overhead irrigation of foliage. Do not allow public use of treated area during application.

FOR FOLIAR BROADCAST SPRAYS

Mix TriStar[®] 8.5 SL with sufficient water and apply as a foliar spray to obtain thorough and uniform spray coverage of the plants. Choose a finished spray volume appropriate for the size of the plants and amount of foliage which will provide thorough coverage throughout the canopy. Apply as soon as insects reach treatment thresholds.

TriStar 8.5 SL mixes quickly in water. This product has been found to be compatible with many commonly used surfactants, miticides and insecticides. Check physical compatibility using the correct proportion of products when combining products without prior history of use.

See label for complete directions for use.

PRECAUTIONS AND RESTRICTIONS

DO NOT make more than 4 applications per year.

DO NOT reapply more than once every 7 days. To determine if application is necessary, monitor pest densities. Consult local extension experts for thresholds.

DO NOT apply more than 25.3 ounces of TriStar 8.5 SL per acre in a single application.

DO NOT apply more than 92.5 ounces of TriStar 8.5 SL per acre per year.

DO NOT apply to bearing fruit trees.

See label for complete application precautions and restrictions.



www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.

©2015 Nufarm. Important: Always read and follow label instructions. TriStar[®] is a trademark of Nippon Soda Co., Ltd. 15-T0-0050