

NEW LOW VOC, WATER-BASED FORMULA

QUICK AND EASY MITE AND LEAFMINER CONTROL.

Minx[™] 2 Insecticide delivers quick and easy broad-spectrum mite and leafminer control in a new low volatile organic compound (VOC), water-based formula. It also works to suppress aphids, whiteflies and thrips. Compared to many other abamectin products, Minx 2 offers added application flexibility and applicator comfort.

COMPETITIVE ADVANTAGES

QUICK CONTROL, EASY ON PLANTS

- New low VOC, water-based formula of abamectin
- Qualifies as low VOC under California regulations
- Contact and systemic control
- Mite and leafminer control
- Suppression of aphids, whiteflies and thrips
- Excellent plant safety

AT-A-GLANCE INFO

EPA REG. NO. 228-736

ACTIVE INGR. abamectin (2.0%)

FORMULATION microemulsion

CHEM. FAMILY avermectin

IRAC NO. 6

SIGNAL WORD caution

RESTRICTED no

PACKAGE SIZE 4 x1qt, 4 x1gal

LANDSCAPE ORNAMENTALS

FIELD-GROWN NURSERY

CONTAINER-GROWN NURSERY

GREENHOUSE



Grow a better tomorrow

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

Safari

SureGuard SC





QUICK AND EASY MITE AND LEAFMINER CONTROL.

KEY USES

Ornamentals

KEY INSECTS CONTROLLED

(See product label for complete list) Aphids Leafminers Mites Thrips Whiteflies

USE SUMMARY

Minx[™] 2 is a microemulsion which when mixed with water according to the directions, will control leafminers and mites and suppress whiteflies, thrips and aphids on ornamentals. Mix with sufficient water and apply as a foliar spray to obtain uniform coverage.

FOR MITES: Apply when mites first appear and repeat as necessary to maintain control.

FOR LEAFMINERS: Apply as needed and repeat at 7-day intervals or as necessary to maintain control.

FOR SUPPRESSION OF APHIDS, THRIPS AND WHITEFLIES: Apply when young, immature stages of these pests are first observed and repeat every 7 days for two or three weeks. After which time, rotate to other products that have different modes of action than Minx 2 for at least two to three weeks. Aphids, thrips and whiteflies are killed by direct contact with the Minx 2 spray. Residual control of pests may be enhanced with the addition of a horticultural spray oil at 0.5 to 1.0% of the spray volume on woody ornamentals and landscape plants. Some plants are sensitive to oils, the user should spray a small number of plants and observe plants for two weeks before spraying the remaining plants. Repeat application as necessary, but no sooner than 7 days to maintain control. Excessive cold or warm temperatures may increase the chance of plant damage following application with oils.

PROVEN PERFORMANCE

RATE	PESTS
4 fl oz/100 gal	Mites: European red mite, two-spotted spider mite, carmine spider mite, southern red mite, spruce spider mite
	Tarsonemid mites: Cyclamen and broad mites
	Eriophyid mites: Rust and bud mites
8 fl oz/100 gal	Aphids, thrips and whiteflies
	Leafminers
	Boxwood leafminer

See label for complete directions for use.

PRECAUTIONS AND RESTRICTIONS

Do not use Minx 2 for suppression of aphids, whiteflies and thrips on roses, chrysanthemums and gerbera. These ornamentals are primary hosts of mites and Liriomyza leafminers for which Minx 2 applications should be targeted. Additional applications of this product to suppress aphids, thrips and whiteflies on these plants will increase the selection pressure on mites and leafminers which may result in greater tolerance to Minx 2 among these pests.

See label for complete application precautions and restrictions.



nufarm.com/usturf



For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/usturf to download a full label.

©2019 Nufarm. Important: Always read and follow label instruction. 3336® is a trademark of Nufarm. Minx™ is trademark of Cleary Chemical, LLC. Safari® and SureGuard® are trademarks of Valent U.S.A. LLC